

# Spatial use of an agrarian landscape by linnets (Linaria cannabina) in late

summer in Central Europe

Carlos Gutiérrez-Expósito<sup>1</sup>, Benedikt Giessing<sup>1</sup>, Jörg Hahne<sup>2</sup>, Carmen Vives-Ferrándiz<sup>1</sup>, Christian Wolf<sup>1</sup>





## STUDY GOALS

- Assess the daily activity pattern and habitat use by linnets
- Identify the habitat preferences of linnets in a farmland landscape in late summer




Figure 1. Example of the positions of one of the tracked Linnet and the resulting Minimum Convex Polygon (MCP)




Figure 2. Proportion of time spent in the different habitats

#### **METHODS**

## Radio-tracking

- 21 linnets were equipped with BioTrack PicoPip tags
- Continuous tracking for one entire day

## Analysis

- Time spent in each habitat
- Proportion of habitat available in the home range
- Jacobs' habitat selectivity index
- Significance by paired samples Wilcoxon test



STUDY AREA




Figure 3. Linnet's habitat Jacobs' selectivity index (+1 = highly selected; -1 = completely avoided)

## CONCLUSIONS

- O No differences in time spent in arboreal vs. herbaceous habitats (W=188 p<0.05)
- Among arboreal habitats, tree patches were preferred over human settlement (W=460, p<0.005)
- O Harvested fields were highly selected, while drilled or growing crops were avoided (W=57, p<0.005)



<sup>1</sup>tier3 solutions GmbH, Kolberger Str. 61-63, Leverkusen, Germany, e-mail: <a href="mailto:christian.wolf@tier3.de">christian.wolf@tier3.de</a>

<sup>2</sup> Bayer AG – Crop Science Division, BAG-CS-RD-Rs-EnSa Alfred Nobel Str. 50, 40789 Monheim am Rhein, Germany